

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

CERTIFICATE OF MOLD ANALYSIS

PREPARED FOR:

SANDUSKY BAY INSPECTIONS LLC

PHONE NUMBER: (419) 503-3644

EMAIL: INFO@SANDUSKYBAYINSPECTIONS.COM

TEST LOCATION:

SAMPLE CLIENT

1234 SAMPLE ST

SANDUSKY, OH 44870

CHAIN OF CUSTODY #: 52034724

COLLECTED: WED MAY 24, 2017

RECEIVED: THU MAY 25, 2017

REPORTED: THU MAY 25, 2017



**APPROVED BY: JOHN D. SHANE PH.D.,
LABORATORY MANAGER**

VERSION: 1.0 (A VERSION NUMBER GREATER THAN ONE (1) INDICATES THAT THE DATA IN THIS REPORT HAS BEEN AMENDED)

EPA regulations or standards for airborne or surface mold concentrations have not been established. There are also no EPA regulations or standards for evaluating health effects due to mold exposure. Information about mold can be found at www.epa.gov/mold.

All samples were received in an acceptable condition for analysis unless noted specifically in the Comments section under a particular sample. All results relate only to the samples submitted for analysis.

A version greater than 1.0 indicates that the lab report has been revised.

FOR MORE INFORMATION, PLEASE CONTACT INSPECTORLAB AT (888) 854-0477 OR EMAIL ASK@INSPECTORLAB.COM

Detailed Mold Report (NAMES IN RED ARE WATER-INDICATING FUNGI)

Analysis Method	Air Analysis	Air Analysis	Air Analysis	Surface Analysis
Lab Sample #	52034724-1	52034724-2	52034724-3	52034724-4
Sample Identification	21023444	21045322	21042361	SWAB #2
Sample Location	OUTSIDE CONTROL	MASTER BEDROOM	TV ROOM	TV ROOM CLOSET WALL
Sample Type / Metric	Air-O-Cell/150.0L	Air-O-Cell/150.0L	Air-O-Cell/150.0L	Air-O-Cell
Analysis Date	Wed June 10, 2015			
Determination	CONTROL	NORMAL	PROBLEM	GROWTH

Fungal Types Identified	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total	Mold Present	
*INDOOR PROBLEM FUNGI											
Cladosporium sphaerospermum	---	---	---	---	---	---	81	543	13	X	
Penicillium	---	---	---	---	---	---	---	---	---	X	
Penicillium/Aspergillus	---	---	---	---	---	---	472	3,162	80	---	
**Non-Problem Fungi											
Alternaria	7	47	2	---	---	---	3	20	<1	---	
Ascospores	19	127	7	17	114	9	---	---	---	---	
Basidiospores	47	315	17	7	47	4	9	60	1	---	
Bipolaris/Drechslera	5	34	1	1	7	<1	---	---	---	---	
Cladosporium	124	831	47	37	248	21	3	20	<1	---	
Curvularia	7	47	2	1	7	<1	---	---	---	---	
Epicoccum	6	40	2	2	13	1	---	---	---	---	
Nigrospora	4	27	1	---	---	---	---	---	---	---	
Penicillium/Aspergillus	6	40	2	91	610	52	---	---	---	---	
Pithomyces	6	40	2	2	13	1	---	---	---	---	
Smut/Myxomycetes	31	208	11	14	94	8	17	114	2	---	
Total Spore Count	262	1,756	100	172	1,153	100	585	3,919	100	N/A	
Minimum Detection Limit	7			7			7			N/A	
Background Debris											
Minerals	---	---	---	---	---	---	12	80	50	---	
Newspaper Ink	---	---	---	---	---	---	3	20	12	---	
Oil Droplets	---	---	---	---	---	---	9	60	37	---	
Comments/Definitions	CONTROL samples are normally taken outside a building to provide a baseline from which samples on the interior of the building are compared. Outside air is considered normal whatever the mold counts may be. Light Debris: The debris present in the sample likely had no effect on the accuracy of the mold count.			Mold counts are within a NORMAL RANGE and there is no indication, based on the mold counts, that there is any exposure concern to the occupants. The LIGHT DEBRIS present in the sample likely had no effect on the accuracy of the mold count.			Mold concentrations in the air are ABNORMAL and based on the mold counts, you likely have a mold source from which spores are able to become airborne and are an exposure concern to the occupants. MODERATE DEBRIS: in the sample likely had limited affect on the accuracy of the mold count.			Presence of current or former MOLD GROWTH observed. EXPOSURE TO SPORES LIKELY and will continue if the growth is not removed. An active or intermittent water source will cause the mold to continue to grow if the water source is not eliminated.	
Raw Count:	Actual number of spores observed and counted.										
Spores/m³:	Spores per cubic meter.										
% of Total:	Percentage of a particular spore in relation to total number of spores.										
X:	Spore type was observed.										
--- :	Spore type was not observed.										

* Indoor Problem Fungi are generally capable of growing on wetted building materials.
 ** Non-Problem Fungi are less capable or do not grow on wetted building materials. They are commonly found in the air outside and infiltrate into indoor air naturally. High numbers of any one of these spore types as compared to the Control sample may indicate that they are growing on wetted building materials indoors.
 Spore types not listed in this report were not observed.
 Background debris estimates the amount of non-spore particles. Increasing amount of debris will affect the accuracy of the spore counts. Total percent may not equal 100% due to rounding.

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

Introduction

All spores found in indoor air are also normally found in outdoor air because most originate or live in the soil and on dead or decaying plants. Therefore, it is not unusual to find mold spores in indoor air. This Mold Glossary is only intended to provide general information about the mold found in the samples that were provided to the laboratory.

Alternaria

Outdoor Habitat: One of the most commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted.

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis), Common cause of extrinsic asthma

Disease Potential: Not normally considered a pathogen, but can become so in immunocompromised persons.

Toxin Potential: Several known

Comments: One of the most common and potent allergens in the indoor and outdoor air. Seen in indoor air in low concentrations, probably as a result of outdoor air infiltration and/or recycling of settled dust.

Ascospores

Outdoor Habitat: Soil and decaying vegetation, dead and dying insects. These spores constitute a large part of the spores in the air and can be found in the air in very large numbers in the spring and summer, especially during and up to three (3) days after a rain.

Indoor Habitat: Very few of fungi that produce ascospores grow indoors. Some fungi that produce ascospores are recognizable by their spores and when observed are listed under their own categories. Wetted wood and gypsum wallboard paper

Allergy Potential: Depends on the type of fungus producing the ascospores.

Disease Potential: Not normally pathogenic as a group

Toxin Potential: None known

Comments: Ascospores are produced from a very large group of fungi. Notable ascospores that are considered problematic for indoor environments are Chaetomium, Peziza, and Ascotracha. If these types of ascspores are observed they will be listed in the report under their own names.

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

Basidiospores

Outdoor Habitat: These are mushroom spores and are common everywhere, especially in the late summer and fall.

Indoor Habitat: Very wet wood products, especially on footer plates, basements, and crawlspaces. Sometimes mushrooms can be observed growing in potted plants indoors.

Allergy Potential: Rarely reported, but some Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis) has been reported.

Disease Potential: None known

Toxin Potential: None known

Comments: This group includes wood rotting fungi, including dry rot (*Serpula* and *Poria*) that are especially destructive to buildings. However, if these types of spores (dry rot group) are observed in the sample they are listed under their own names on the report.

Bipolaris/Drechslera

Outdoor Habitat: Commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: None known

Comments: This category represents at least three genera, including *Bipolaris*, *Drechslera*, and *Exserohilum*. This group cannot be consistently separated by spore morphology alone.

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

Cladosporium

- Outdoor Habitat:** Cladosporium is one of the most common environmental fungi observed worldwide. Soil and decaying vegetation. Cladosporium herbarum and C. cladosporioides are among the most frequently encountered species, both in outdoor and indoor environments
- Indoor Habitat:** Wetted wood and gypsum wallboard paper, paper products, textiles, rubber, window sills
- Allergy Potential:** Type I (hay fever, asthma) - an important and common outdoor allergen
- Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals. Cladosporium are some of the most common species reported as indoor contaminants, occasionally linked to health problems.
- Toxin Potential:** Two known, but not highly toxic
- Comments:** The most commonly reported spore in the outdoor air worldwide. An important and common allergen source.
-

Cladosporium sphaerospermum

- Outdoor Habitat:** Dead or dying cellulosic materials like wood and leaves.
- Indoor Habitat:** A favorite place for this mold type to grow is on wetted or rotting window sills. Also likes to grow on all type of wetted or rotting wood.
- Allergy Potential:** Type I (hay fever, asthma) - an important and common outdoor allergen
- Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.
- Toxin Potential:** None known
- Comments:** A common mold on wetted wood, especially on window sills. Not frequently found in the air.
-

Curvularia

- Outdoor Habitat:** Soil and decaying vegetation
- Indoor Habitat:** Wetted wood and gypsum wallboard paper, many cellulytic substrates
- Allergy Potential:** Type I (hay fever, asthma), common cause of allergenic rhinitis
- Disease Potential:** Potential human pathogen in immunocompromised people
- Toxin Potential:** None known
- Comments:** None
-

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

Epicoccum

Outdoor Habitat: Epicoccum is a widespread cosmopolitan that grows on dead or decaying organic matter, wood, textiles, paper, a variety of foods, insects and human skin. It is commonly found in the soil. Epicoccum spores are more prevalent on dry, windy days, with higher counts late in the day.

Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted such as gypsum board, floors, carpets, mattress dust, and house plants.

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: None known

Toxin Potential: None known

Comments: Very common in outdoor air in the summer months, especially in the midwest USA during harvest times.

Nigrospora

Outdoor Habitat: Soil and decaying vegetation

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: None known

Toxin Potential: None known

Comments: Rarely observed growing indoors

Penicillium

Outdoor Habitat: Soil and decaying vegetation, textiles, fruits

Indoor Habitat: Wetted wood and gypsum wallboard paper, textiles, leather

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Several known

Comments: Extremely common in indoor air, but without the fruiting bodies associated with the spores will be listed as "Penicillium / Aspergillus" group. Penicillium identified in air samples indicates that the fruiting bodies were observed. This usually suggests that the source of the mold is nearby and / or a growth was disturbed. The fruiting bodies are not easily sent airborne, nor do they stay in the air long.

PREPARED FOR: SANDUSKY BAY INSPECTIONS LLC

TEST ADDRESS: 1234 SAMPLE ST SANDUSKY, OH 44870

Penicillium/Aspergillus

Outdoor Habitat: Soil and decaying vegetation, textiles, fruits. These spores are commonly observed and are a normal part of outside air.

Indoor Habitat: Wetted wood and gypsum wallboard paper, textiles, leather, able to grow on many types of substrates.

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Several known

Comments: Extremely common in indoor air in low amounts. This type of spore should not constitute an overwhelming percentage and be present in very high numbers.

These two genera are grouped together because they cannot be reliably differentiated into their respective genera based solely on spore morphology.

Pithomyces

Outdoor Habitat: Soil and decaying vegetation and their spores are easily dispersed into the air by wind

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: None known

Disease Potential: None known

Toxin Potential: One known (sporidesmin)

Comments: A very common spore type in the air. Can be a water indicator mold type indoors

Smut/Myxomycetes

Outdoor Habitat: Soil and decaying vegetation and wood, especially dead stumps and bark

Indoor Habitat: Not known to grow indoors, sometimes found on firewood

Allergy Potential: Type I (hay fever, asthma), rare

Disease Potential: None known

Toxin Potential: None known

Comments: These two groups are difficult to distinguish due to their "round, brown" morphology. Smuts are especially common in the environment and can be seen in indoor air samples even during the winter in homes because the spores can get trapped in carpets
